

*MODULE 5*

# APPROVAL OR DENIAL

Coordinator Corner has developed this summary to assist your learning and help you fully understand the information featured in this video

## Video breakdown by topic

### **1 Topic: What to do when a referral is approved** | Time: 0:15

Once you have been successful in getting a Synagis® (palivizumab) patient referral approved, there is still work to be done. After receiving notification of an approved referral:

- Promptly coordinate with parents to schedule the high-risk baby's first, or next, Synagis dose
  - If a home care agency or other clinic will be responsible for dosing, make sure to communicate with them to ensure a dosing schedule is set
  - To begin scheduling dosing:
    - Confirm if the baby already received his or her first dose of Synagis while in the NICU or the nursery
      - If he or she hasn't, and if RSV season hasn't started yet, find out when dosing can start in your area by checking with a MedImmune representative or the RSVAlert® website
    - If RSV season is already in full swing and the baby has not received the first dose, schedule it as soon as possible
    - If the baby has already received the first dose, schedule the next one 28 to 30 days later

### **INDICATION & SELECT SAFETY INFORMATION**

Synagis is indicated for the prevention of serious lower respiratory tract disease caused by respiratory syncytial virus (RSV) in children at high risk of RSV disease.

The recommended dose of Synagis is 15 mg/kg of body weight given monthly by intramuscular injection. The first dose of Synagis should be administered prior to commencement of the RSV season and the remaining doses should be administered monthly throughout the RSV season.

**Please see Important Safety Information on page 4, and accompanying full Prescribing Information for Synagis, including Patient Information.**

**2 Topic: Setting expectations with parents and caregivers | Time: 1:14**

- After the first dose has been scheduled, parents and caregivers may not know what to expect next. Here's how you can help them:
  - Explain that they should schedule all remaining Synagis® (palivizumab) appointments at once, each between 28 and 30 days apart, so future scheduling won't get overlooked
  - Make sure to reinforce the importance of timely dosing
  - Clarify that Synagis is NOT a vaccine, but a medication that provides enough antibodies to help protect their baby against severe RSV disease for 1 month
- Talk to parents about the following:
  - Insurance coverage for Synagis and any potential out-of-pocket costs for which they may be responsible
  - Access 360™, a Synagis support program, which may be able to provide financial assistance
    - Eligible parents can submit a claim to be reimbursed for the cost of Synagis and/or administration, by filling out an Access 360 claim form
  - Potential consent-to-ship phone calls from the Specialty Pharmacy Provider (SPP), so these calls do not come as a surprise
  - Potential scheduling calls from a home health nurse (if Synagis will be administered by a home health agency)
- Coordinate With SPP for medication supply
  - Confirm how you will be receiving communication from the SPP (e.g., fax, email)
  - Communicate often with the SPP to ensure Synagis is delivered on time and in proper dosage
  - Periodically verify patient's information with the SPP
    - *Note: Access 360 can also follow up with the SPP*

**3 Topic: What to do when a referral is denied | Time: 3:04**

- When a referral is denied, remember that there are still options available to appeal the insurance company's decision. Upon receiving denial letter from the payer:
  - Identify the exact reason for denial
    - Clinical (e.g., patient not being within payer policy)
    - Administrative (e.g., a missing signature or ICD-9 or ICD-10 code)
  - Request a written denial and keep a record of the name and contact information of the person who handles your request
    - You will also want to pinpoint the necessary steps for appealing the denial, which can usually be found in the denial letter
      - Take care to follow the appeals process exactly as outlined. If the letter does not state these steps, contact the payer directly to find out what they are
  - Discuss referral denial with parents/caregivers
    - Parents may feel discouraged and unsure of what to do next; you'll want to be reassuring and keep them informed about possibilities and next steps
    - Encourage them to be proactive by contacting their insurance provider and/or their employer to find out what options they may have
    - Educate parents on how they may counteract the denial by appealing it

**Please see Important Safety Information on page 4, and accompanying full Prescribing Information for Synagis, including Patient Information.**



**4 Topic: Appealing the denial | Time: 4:32**


- The earlier you file the appeal, the better chance you have of preventing unnecessary holdups
  - If you have multiple appeals in your queue, group them together to streamline the process
  - Contact payer within 24 hours
  - Keep parents informed throughout the appeals process
- Preparing an appeal letter
  - List all of the risk factors and clinically relevant information that puts the patient at high risk for severe RSV disease. This may include:
    - Local RSV virology data (RSVAlert.com) if you are filing the appeal during the RSV season
    - Letter from a medical specialist, such as a pulmonologist, cardiologist, or neonatologist

**5 Topic: When an appeal does not overturn a denial | Time: 5:42**

- When an appeal doesn't result in an overturned denial, you still have a few options for disputing the denial
  - Request a peer-to-peer review with a medical director
  - Request a third-party review
    - The insurance company will consult with an external specialist, such as a cardiologist or pulmonologist
  - In peer-to-peer and third-party reviews, the medical professionals may be able to identify potential errors in the reasons given for the denial, as well as provide further support for the medical necessity of Synagis® (palivizumab)

**6 Topic: Access 360 appeal support | Time: 6:20**

- Access 360 cannot actually submit an appeal on your behalf, but it CAN provide you with:
  - Payer-specific forms, instructions, and contact information
  - Follow up communications with the payer on the outcome of the appeal
  - *Note: The AstraZeneca Medical Information team can also provide you with additional clinical supporting documentation*
- If your appeal is unsuccessful, Access 360 offers reimbursement assistance options to parents through its various support programs
  - Contact your MedImmune field reimbursement manager (FRM) for denial-specific support if there is a signed Patient Authorization Form (PAF) on file



If you want to become an official certified Synagis coordinator, Coordinator Corner will help you every step of the way. All you have to do is view all the modules and take all the checkpoints to become certified and eligible to receive an educational textbook, or you can make a charitable donation.

You must be registered to be eligible for certification—**so register today.**

**SELECT SAFETY INFORMATION**

Adverse reactions occurring greater than or equal to 10% and at least 1% more frequently than placebo are fever and rash. In post-marketing reports, cases of severe thrombocytopenia (platelet count <50,000/microliter) and injection site reactions have been reported.

**Please see Important Safety Information on page 4, and accompanying full Prescribing Information for Synagis, including Patient Information.**

## IMPORTANT SAFETY INFORMATION

Synagis® (palivizumab) is indicated for the prevention of serious lower respiratory tract disease caused by respiratory syncytial virus (RSV) in children at high risk of RSV disease. Safety and efficacy were established in children with bronchopulmonary dysplasia (BPD), infants with a history of premature birth ( $\leq 35$  weeks gestational age), and children with hemodynamically significant congenital heart disease (CHD). The recommended dose of Synagis is 15 mg/kg of body weight given monthly by intramuscular injection. The first dose of Synagis should be administered prior to commencement of the RSV season and the remaining doses should be administered monthly throughout the RSV season. Children who develop an RSV infection should continue to receive monthly doses throughout the RSV season.

The efficacy of Synagis at doses less than 15 mg/kg, or of dosing less frequently than monthly throughout the RSV season, has not been established.

Synagis is contraindicated in children who have had a previous significant hypersensitivity reaction to Synagis. Cases of anaphylaxis and anaphylactic shock, including fatal cases, have been reported following initial exposure or re-exposure to Synagis. Other acute hypersensitivity reactions, which may be severe, have also been reported on initial exposure or re-exposure to Synagis. The relationship between these reactions and the development of antibodies to Synagis is unknown. If a significant hypersensitivity reaction occurs with Synagis, its use should be permanently discontinued. If a mild hypersensitivity reaction occurs, clinical judgment should be used regarding cautious readministration of Synagis. As with any intramuscular injection, Synagis should be given with caution to children with thrombocytopenia or any coagulation disorder. Palivizumab may interfere with immunological-based RSV diagnostic tests, such as some antigen detection-based assays.

Adverse reactions occurring greater than or equal to 10% and at least 1% more frequently than placebo are fever and rash. In post-marketing reports, cases of severe thrombocytopenia (platelet count  $< 50,000/\text{microliter}$ ) and injection site reactions have been reported.

**Please see accompanying full Prescribing Information for Synagis, including Patient Information.**



## HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use SYNAGIS safely and effectively. See full prescribing information for SYNAGIS.

**SYNAGIS® (palivizumab) injection, for intramuscular use**  
**Initial U.S. Approval: 1998**

### INDICATIONS AND USAGE

Synagis is a respiratory syncytial virus (RSV) F protein inhibitor monoclonal antibody indicated for the prevention of serious lower respiratory tract disease caused by RSV in children at high risk of RSV disease.

- Safety and efficacy were established in children with bronchopulmonary dysplasia (BPD), infants with a history of premature birth (less than or equal to 35 weeks gestational age), and children with hemodynamically significant congenital heart disease (CHD).
- The safety and efficacy of Synagis have not been established for treatment of RSV disease. (1)

### DOSAGE AND ADMINISTRATION

15 mg per kg of body weight, administered intramuscularly prior to commencement of the RSV season and remaining doses administered monthly throughout the RSV season. (2.1)

Children undergoing cardio-pulmonary bypass should receive an additional dose of Synagis as soon as possible after the cardio-pulmonary bypass procedure (even if sooner than a month from the previous dose). Thereafter, doses should be administered monthly as scheduled. (2.1, 12.3)

### DOSAGE FORMS AND STRENGTHS

Single-dose liquid solution vials: 50 mg per 0.5 mL and 100 mg per 1 mL. (3)

## FULL PRESCRIBING INFORMATION: CONTENTS\*

- 1 INDICATIONS AND USAGE
- 2 DOSAGE AND ADMINISTRATION
  - 2.1 Dosing Information
  - 2.2 Administration Instructions
- 3 DOSAGE FORMS AND STRENGTHS
- 4 CONTRAINDICATIONS
- 5 WARNINGS AND PRECAUTIONS
  - 5.1 Hypersensitivity Reactions
  - 5.2 Coagulation Disorders
  - 5.3 RSV Diagnostic Test Interference
  - 5.4 Treatment of RSV Disease
  - 5.5 Proper Administration
- 6 ADVERSE REACTIONS
  - 6.1 Clinical Studies Experience
  - 6.2 Postmarketing Experience

## FULL PRESCRIBING INFORMATION

### 1 INDICATIONS AND USAGE

Synagis is indicated for the prevention of serious lower respiratory tract disease caused by respiratory syncytial virus (RSV) in children at high risk of RSV disease.

*The following points should be considered when prescribing Synagis:*

- Safety and efficacy were established in children with bronchopulmonary dysplasia (BPD), infants with a history of premature birth (less than or equal to 35 weeks gestational age), and children with hemodynamically significant congenital heart disease (CHD) [see *Clinical Studies* (14)].
- The safety and efficacy of Synagis have not been established for treatment of RSV disease.

### 2 DOSAGE AND ADMINISTRATION

#### 2.1 Dosing Information

The recommended dose of Synagis is 15 mg per kg of body weight given monthly by intramuscular injection. The first dose of Synagis should be administered prior to commencement of the RSV season and the remaining doses should be administered monthly throughout the RSV season. Children who develop an RSV infection should continue to receive monthly doses throughout the RSV season. In the northern hemisphere, the RSV season typically commences in November and lasts through April, but it may begin earlier or persist later in certain communities.

Synagis serum levels are decreased after cardio-pulmonary bypass [see *Clinical Pharmacology* (12.3)]. Children undergoing cardio-pulmonary bypass should receive an additional dose of Synagis as soon as possible after the cardio-pulmonary bypass procedure (even if sooner than a month from the previous dose). Thereafter, doses should be administered monthly as scheduled.

The efficacy of Synagis at doses less than 15 mg per kg, or of dosing less frequently than monthly throughout the RSV season, has not been established.

#### 2.2 Administration Instructions

- **DO NOT DILUTE THE PRODUCT.**
- **DO NOT SHAKE OR VIGOROUSLY AGITATE THE VIAL.**
- Parenteral drug products should be inspected visually for particulate matter and discoloration prior to administration. Do not use any vials exhibiting particulate matter or discoloration.

### CONTRAINDICATIONS

Previous significant hypersensitivity reaction to Synagis. (4)

### WARNINGS AND PRECAUTIONS

- Anaphylaxis and anaphylactic shock (including fatal cases), and other severe acute hypersensitivity reactions have been reported. Permanently discontinue Synagis and administer appropriate medications if such reactions occur. (5.1)
- As with any intramuscular injection, Synagis should be given with caution to children with thrombocytopenia or any coagulation disorder. (5.2)
- Palivizumab may interfere with immunological-based RSV diagnostic tests such as some antigen detection-based assays. (5.3, 12.4)

### ADVERSE REACTIONS

Adverse reactions occurring greater than or equal to 10% and at least 1% more frequently than placebo are fever and rash. (6.1)

**To report SUSPECTED ADVERSE REACTIONS, contact MedImmune at 1-877-633-4411 or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).**

### USE IN SPECIFIC POPULATIONS

Safety and effectiveness in children greater than 24 months of age at the start of dosing have not been established. (8.4)

See 17 for PATIENT COUNSELING INFORMATION and FDA-approved patient labeling.

Revised: 03/2014

### 7 DRUG INTERACTIONS

### 8 USE IN SPECIFIC POPULATIONS

- 8.1 Pregnancy
- 8.4 Pediatric Use

### 10 OVERDOSAGE

### 11 DESCRIPTION

### 12 CLINICAL PHARMACOLOGY

- 12.1 Mechanism of Action
- 12.3 Pharmacokinetics
- 12.4 Microbiology

### 13 NONCLINICAL TOXICOLOGY

- 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

### 14 CLINICAL STUDIES

### 16 HOW SUPPLIED/STORAGE AND HANDLING

### 17 PATIENT COUNSELING INFORMATION

\*Sections or subsections omitted from the full prescribing information are not listed.

- Using aseptic techniques, attach a sterile needle to a sterile syringe. Remove the flip top from the Synagis vial and wipe the rubber stopper with a disinfectant (e.g., 70% isopropyl alcohol). Insert the needle into the vial and withdraw into the syringe an appropriate volume of solution. Administer immediately after drawing the dose into the syringe.
- Synagis should be administered in a dose of 15 mg per kg intramuscularly using aseptic technique, preferably in the anterolateral aspect of the thigh. The gluteal muscle should not be used routinely as an injection site because of the risk of damage to the sciatic nerve. The dose (volume of injection in mL) per month = patient weight (kg) x 15 mg per kg ÷ 100 mg per mL of Synagis. Injection volumes over 1 mL should be given as a divided dose.
- Synagis is supplied as a single-dose vial and does not contain preservatives. Do not re-enter the vial after withdrawal of drug; discard unused portion. Only administer one dose per vial.
- Use sterile disposable syringes and needles. To prevent the transmission of hepatitis viruses or other infectious agents from one person to another, DO NOT reuse syringes and needles.

### 3 DOSAGE FORMS AND STRENGTHS

Single-dose liquid solution vials: 50 mg per 0.5 mL and 100 mg per 1 mL.

### 4 CONTRAINDICATIONS

Synagis is contraindicated in children who have had a previous significant hypersensitivity reaction to Synagis [see *Warnings and Precautions* (5.1)].

### 5 WARNINGS AND PRECAUTIONS

#### 5.1 Hypersensitivity Reactions

Cases of anaphylaxis and anaphylactic shock, including fatal cases, have been reported following initial exposure or re-exposure to Synagis. Other acute hypersensitivity reactions, which may be severe, have also been reported on initial exposure or re-exposure to Synagis. Signs and symptoms may include urticaria, pruritus, angioedema, dyspnea, respiratory failure, cyanosis, hypotonia, hypotension, and unresponsiveness. The relationship between these reactions and the development of antibodies to Synagis is unknown. If a significant hypersensitivity reaction occurs with Synagis, its use should be permanently discontinued. **If anaphylaxis or other significant hypersensitivity reaction occurs, administer appropriate medications (e.g., epinephrine) and provide supportive care as required.** If a mild hypersensitivity reaction occurs, clinical judgment should be used regarding cautious readministration of Synagis.

## 5.2 Coagulation Disorders

Synagis is for intramuscular use only. As with any intramuscular injection, Synagis should be given with caution to children with thrombocytopenia or any coagulation disorder.

## 5.3 RSV Diagnostic Test Interference

Palivizumab may interfere with immunological-based RSV diagnostic tests such as some antigen detection-based assays. In addition, palivizumab inhibits virus replication in cell culture, and therefore may also interfere with viral culture assays. Palivizumab does not interfere with reverse transcriptase-polymerase chain reaction based assays. Assay interference could lead to false-negative RSV diagnostic test results. Therefore, diagnostic test results, when obtained, should be used in conjunction with clinical findings to guide medical decisions [see *Microbiology* (12.4)].

## 5.4 Treatment of RSV Disease

The safety and efficacy of Synagis have not been established for treatment of RSV disease.

## 5.5 Proper Administration

The single-dose vial of Synagis does not contain a preservative. Administration of Synagis should occur immediately after dose withdrawal from the vial. The vial should not be re-entered. Discard any unused portion.

## 6 ADVERSE REACTIONS

The most serious adverse reactions occurring with Synagis are anaphylaxis and other acute hypersensitivity reactions [see *Warnings and Precautions* (5.1)].

### 6.1 Clinical Studies Experience

*Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.*

The data described below reflect exposure to Synagis (n=1639) compared with placebo (n=1143) in children 3 days to 24.1 months of age at high risk of RSV-related hospitalization in two clinical trials. Trial 1 was conducted during a single RSV season and studied a total of 1502 children less than or equal to 24 months of age with BPD or infants with premature birth (less than or equal to 35 weeks gestation) who were less than or equal to 6 months of age at study entry. Trial 2 was conducted over four consecutive seasons among a total of 1287 children less than or equal to 24 months of age with hemodynamically significant congenital heart disease.

In Trials 1 and 2 combined, fever and rash were each reported more frequently among Synagis than placebo recipients, 27% versus 25%, and 12% versus 10%, respectively. Adverse reactions observed in the 153-patient crossover study comparing the liquid and lyophilized formulations were comparable for the two formulations, and were similar to those observed with Synagis in Trials 1 and 2.

#### *Immunogenicity*

In Trial 1, the incidence of anti-palivizumab antibody following the fourth injection was 1.1% in the placebo group and 0.7% in the Synagis group. In children receiving Synagis for a second season, one of the fifty-six children had transient, low titer reactivity. This reactivity was not associated with adverse events or alteration in serum concentrations. Immunogenicity was not assessed in Trial 2.

A trial of high-risk preterm children less than or equal to 24 months of age was conducted to evaluate the immunogenicity of the lyophilized formulation of Synagis (used in Trials 1 and 2 above) and the liquid formulation of Synagis. Three hundred seventy-nine children contributed to the 4 to 6 months post-final dose analysis. The rate of anti-palivizumab antibodies at this time point was low in both formulation groups (anti-palivizumab antibodies were not detected in any subject in the liquid formulation group and were detected in one subject in the lyophilized group (0.5%), with an overall rate of 0.3% for both treatment groups combined).

These data reflect the percentage of children whose test results were considered positive for antibodies to palivizumab in an enzyme-linked immunosorbent assay (ELISA) and are highly dependent on the sensitivity and specificity of the assay.

The ELISA has substantial limitations in detecting anti-palivizumab antibodies in the presence of palivizumab. Immunogenicity samples tested with the ELISA assay likely contained palivizumab at levels that may interfere with the detection of anti-palivizumab antibodies.

An electrochemical luminescence (ECL) based immunogenicity assay, with a higher tolerance for palivizumab presence compared to the ELISA, was used to evaluate the presence of anti-palivizumab antibodies in subject samples from two additional clinical trials. The rates of anti-palivizumab antibody positive results in these trials were 1.1% and 1.5%.

### 6.2 Postmarketing Experience

The following adverse reactions have been identified during post approval use of Synagis. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

**Blood and Lymphatic System Disorders:** severe thrombocytopenia (platelet count less than 50,000 per microliter)

**General Disorders and Administration Site Conditions:** injection site reactions

Limited information from post-marketing reports suggests that, within a single RSV season, adverse events after a sixth or greater dose of Synagis are similar in character and frequency to those after the initial five doses.

## 7 DRUG INTERACTIONS

No formal drug-drug interaction studies were conducted. In Trial 1, the proportions of children in the placebo and Synagis groups who received routine childhood vaccines, influenza vaccine, bronchodilators, or corticosteroids were similar and no incremental increase in adverse reactions was observed among children receiving these agents.

## 8 USE IN SPECIFIC POPULATIONS

### 8.1 Pregnancy

Pregnancy Category C: Synagis is not indicated for adult usage. It is not known whether Synagis can cause fetal harm or could affect reproductive capacity when administered to a pregnant woman.

#### *Animal Data*

Animal reproduction studies have not been conducted.

### 8.4 Pediatric Use

The safety and effectiveness of Synagis in children greater than 24 months of age at the start of dosing have not been established.

## 10 OVERDOSAGE

Overdoses with doses up to 85 mg per kg have been reported in clinical studies and post-marketing experience with Synagis, and in some cases, adverse reactions were reported. In case of overdosage, it is recommended that the patient be monitored for any signs or symptoms of adverse reactions and appropriate symptomatic treatment instituted.

## 11 DESCRIPTION

Palivizumab is a humanized monoclonal antibody (IgG1κ) produced by recombinant DNA technology, directed to an epitope in the A antigenic site of the F protein of RSV. Palivizumab is a composite of human (95%) and murine (5%) antibody sequences. The human heavy chain sequence was derived from the constant domains of human IgG1 and the variable framework regions of the V<sub>H</sub> genes Cor and Cess. The human light chain sequence was derived from the constant domain of Cκ and the variable framework regions of the V<sub>L</sub> gene K104 with Jκ-4. The murine sequences were derived from a murine monoclonal antibody, Mab 1129, in a process that involved the grafting of the murine complementarity determining regions into the human antibody frameworks. Palivizumab is composed of two heavy chains and two light chains and has a molecular weight of approximately 148,000 Daltons.

Synagis is supplied as a sterile, preservative-free liquid solution at 100 mg per mL to be administered by intramuscular injection. Thimerosal or other mercury-containing salts are not used in the production of Synagis. The solution has a pH of 6.0 and should appear clear or slightly opalescent.

Each 100 mg single-dose vial of Synagis liquid solution contains 100 mg of palivizumab and also contains chloride (0.5 mg), glycine (0.1 mg), and histidine (3.9 mg), in a volume of 1 mL. Each 50 mg single-dose vial of Synagis liquid solution contains 50 mg of palivizumab and also contains chloride (0.2 mg), glycine (0.06 mg), and histidine (1.9 mg), in a volume of 0.5 mL.

## 12 CLINICAL PHARMACOLOGY

### 12.1 Mechanism of Action

Palivizumab is a recombinant humanized monoclonal antibody with anti-RSV activity [see *Microbiology* (12.4)].

### 12.3 Pharmacokinetics

In children less than or equal to 24 months of age without congenital heart disease (CHD), the mean half-life of palivizumab was 20 days and monthly intramuscular doses of 15 mg per kg achieved mean ± SD 30 day trough serum drug concentrations of 37 ± 21 mcg per mL after the first injection, 57 ± 41 mcg per mL after the second injection, 68 ± 51 mcg per mL after the third injection, and 72 ± 50 mcg per mL after the fourth injection. Trough concentrations following the first and fourth Synagis dose were similar in children with CHD and in non-cardiac patients. In children given Synagis for a second season, the mean ± SD serum concentrations following the first and fourth injections were 61 ± 17 mcg per mL and 86 ± 31 mcg per mL, respectively.

In 139 children less than or equal to 24 months of age with hemodynamically significant CHD who received Synagis and underwent cardio-pulmonary bypass for open-heart surgery, the mean ± SD serum palivizumab concentration was 98 ± 52 mcg per mL before bypass and declined to 41 ± 33 mcg per mL after bypass, a reduction of 58% [see *Dosage and Administration* (2.1)]. The clinical significance of this reduction is unknown.

Specific studies were not conducted to evaluate the effects of demographic parameters on palivizumab systemic exposure. However, no effects of gender, age, body weight, or race on palivizumab serum trough concentrations were observed in a clinical study with 639 children with CHD (less than or equal to 24 months of age) receiving five monthly intramuscular injections of 15 mg per kg of Synagis.

The pharmacokinetics and safety of Synagis liquid solution and Synagis lyophilized formulation administered via intramuscular injection at 15 mg per kg were studied in a cross-over trial of 153 infants less than or equal to 6 months of age with a history of prematurity. The results of this trial indicated that the trough serum concentrations of palivizumab were comparable between the liquid solution and the lyophilized formulation, which was the formulation used in the clinical studies. A population pharmacokinetic analysis was performed across 22 studies in 1800 patients (1684 pediatric and 116 adult patients) to characterize palivizumab pharmacokinetics and inter-subject variability in serum concentrations. Palivizumab pharmacokinetics was described by a two-compartment linear model with an elimination half-life of 24.5 days in pediatric patients. Clearance of palivizumab in a typical pediatric patient (body weight 4.5 kg) less than or equal to 24 months of age without CHD was estimated to be 11 mL per day with a bioavailability of 70% following intramuscular administration. The inter-patient variability in drug clearance was 48.7% (CV%). Covariate analysis did not identify any factors that could account for the inter-patient variability in order to predict serum concentrations a priori in an individual patient.

### 12.4 Microbiology

#### *Mechanism of Action*

Palivizumab, a recombinant humanized monoclonal antibody which provides passive immunity against RSV, acts by binding the RSV envelope fusion protein (RSV F) on the surface of the virus and blocking a critical step in the membrane fusion process. Palivizumab also prevents cell-to-cell fusion of RSV-infected cells.



### Antiviral Activity

The antiviral activity of palivizumab was assessed in a microneutralization assay in which increasing concentrations of antibody were incubated with RSV prior to addition of the human epithelial cells HEP-2. After incubation for 4-5 days, RSV antigen was measured in an ELISA assay. The neutralization titer (50% effective concentration [EC<sub>50</sub>]) is expressed as the antibody concentration required to reduce detection of RSV antigen by 50% compared with untreated virus-infected cells. Palivizumab exhibited median EC<sub>50</sub> values of 0.65 mcg per mL (mean 0.75 ± 0.53 mcg per mL; n=69, range 0.07-2.89 mcg per mL) and 0.28 mcg per mL (mean 0.35 ± 0.23 mcg per mL; n=35, range 0.03-0.88 mcg per mL) against clinical RSV A and RSV B isolates, respectively. The majority of clinical RSV isolates tested (n=96) were collected from subjects across the United States (CA, CO, CT, IL, MA, NC, NY, PA, RI, TN, TX, VA), with the remainder from Japan (n=1), Australia (n=5) and Israel (n=2). These isolates encoded the most common RSV F sequence polymorphisms found among clinical isolates worldwide.

Palivizumab serum concentrations of greater than or equal to 40 mcg per mL have been shown to reduce pulmonary RSV replication in the cotton rat model of RSV infection by 100-fold.

### Resistance

Palivizumab binds a highly conserved region on the extracellular domain of mature RSV F, referred to as antigenic site II or site A, which encompasses amino acids 262 to 275. All RSV mutants that exhibit resistance to palivizumab have been shown to contain amino acid changes in this region on the F protein.

**F protein sequence variations within antigenic site A:** Amino acid substitutions in antigenic site A selected either in cell culture, in animal models, or in human subjects that resulted in palivizumab resistance were N262D, N268I, K272E/M/N/Q/T, and S275F/L. RSV variants expressing the K272N substitution in F protein showed a 5164 ± 1731-fold decrease in susceptibility (i.e., fold increase in EC<sub>50</sub> value) when compared to the wild-type RSV, while variants containing the N262D, S275F/L, or K272E/M/N/Q/T substitutions showed a greater than 25,000-fold decrease in susceptibility to palivizumab. The N268I substitution conferred partial resistance to palivizumab; however, fold changes in susceptibility were not quantified for this mutant. Studies carried out to investigate the mechanism of virus escape from palivizumab showed a correlation between antibody binding and virus neutralization. RSV with substitutions in antigenic site A that were resistant to neutralization by palivizumab did not bind to palivizumab.

At least one of the palivizumab resistance-associated substitutions, N262D, K272E/Q, or S275F/L was identified in 8 of 126 clinical RSV (59 RSV A and 67 RSV B) isolates from subjects who failed immunoprophylaxis, resulting in a combined resistance-associated mutation frequency of 6.3%. A review of clinical findings revealed no association between antigenic A site sequence changes and RSV disease severity among children receiving palivizumab immunoprophylaxis who develop RSV lower respiratory tract disease.

Analysis of 254 clinical RSV isolates (145 RSV A and 109 RSV B) collected from immunoprophylaxis-naïve subjects revealed palivizumab resistance-associated substitutions in 2 (1 with N262D and 1 with S275F), resulting in a resistance-associated mutation frequency of 0.79%.

**F protein sequence variations outside antigenic site A:** In addition to the sequence variations in antigenic site A known to confer palivizumab resistance, F protein substitutions T100A, G139S, N165D/V406I; T326A, V450A in RSV A, and T74I, A147V, I206L, S285G, V450I, T455I in RSV B were identified in viruses isolated from failures of immunoprophylaxis. These substitutions were not identified in RSV F sequences derived from 254 clinical isolates from immunoprophylaxis-naïve subjects and thus are considered treatment-associated and non-polymorphic. Recombinant RSV B encoding the S285G substitution exhibited palivizumab sensitivity (EC<sub>50</sub> value = 0.39 ± 0.02 mcg per mL) similar to recombinant wild-type RSV B (EC<sub>50</sub> value = 0.17 ± 0.02 mcg per mL).

Palivizumab susceptibility of RSV encoding common F protein sequence polymorphisms located proximal to antigenic site A was evaluated. Recombinant RSV A encoding N276S (EC<sub>50</sub> value = 0.72 ± 0.07 mcg per mL), and recombinant RSV B with S276N (EC<sub>50</sub> value = 0.42 ± 0.04 mcg per mL), exhibited sensitivities comparable to the corresponding recombinant wild-type RSV A (EC<sub>50</sub> value = 0.63 ± 0.22 mcg per mL) and RSV B (EC<sub>50</sub> value = 0.23 ± 0.07 mcg per mL). Likewise, RSV B clinical isolates containing the polymorphic variation V278A were at least as sensitive to neutralization by palivizumab (EC<sub>50</sub> range 0.08-0.45 mcg per mL) as laboratory strains of wild-type RSV B (EC<sub>50</sub> value = 0.54 ± 0.08 mcg per mL). No known polymorphic or non-polymorphic sequence variations outside the antigenic site A on RSV F have been demonstrated to render RSV resistant to neutralization by palivizumab.

### Interference of RSV Diagnostic Assays by Palivizumab

Interference with immunologically-based RSV diagnostic assays by palivizumab has been observed in laboratory studies. Rapid chromatographic/enzyme immunoassays (CIA/EIA), immunofluorescence assays (IFA), and direct immunofluorescence assays (DFA) using monoclonal antibodies targeting RSV F protein may be inhibited. Therefore, caution should be used in interpreting negative immunological assay results when clinical observations are consistent with RSV infection. A reverse transcriptase-polymerase chain reaction (RT-PCR) assay, which is not inhibited by palivizumab, may prove useful for laboratory confirmation of RSV infection [see *Warnings and Precautions* (5.3)].

## 13 NONCLINICAL TOXICOLOGY

### 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Carcinogenesis, mutagenesis, and reproductive toxicity studies have not been performed.

## 14 CLINICAL STUDIES

The safety and efficacy of Synagis were assessed in two randomized, double-blind, placebo-controlled trials of prophylaxis against RSV infection in children at high risk of an RSV-related hospitalization. Trial 1 was conducted during a single RSV season and studied a total of 1502 children less than or equal to 24 months of age with BPD or infants with premature birth (less than or equal to 35 weeks gestation) who were less than or equal to 6 months of age at study entry. Trial 2 was conducted over four consecutive seasons among a total of 1287 children less than or equal to 24 months of age with hemodynamically significant congenital heart disease. In both trials participants received 15 mg per kg Synagis or an equivalent volume of placebo via intramuscular injection monthly for five injections and were followed for 150 days from randomization. In Trial 1, 99% of all subjects completed the study and 93% completed all five injections. In Trial 2, 96% of all subjects completed the study and 92% completed all five injections. The incidence of RSV hospitalization is shown in Table 1. The results were shown to be statistically significant using Fisher's exact test.

**Table 1: Incidence of RSV Hospitalization by Treatment Group**

Trial		Placebo	Synagis	Difference Between Groups	Relative Reduction
Trial 1 Impact-RSV	N	500	1002		
	Hospitalization	53 (10.6%)	48 (4.8%)	5.8%	55%
Trial 2 CHD	N	648	639		
	Hospitalization	63 (9.7%)	34 (5.3%)	4.4%	45%

In Trial 1, the reduction of RSV hospitalization was observed both in children with BPD (34/266 [12.8%] placebo versus 39/496 [7.9%] Synagis) and in premature infants without BPD (19/234 [8.1%] placebo versus 9/506 [1.8%] Synagis). In Trial 2, reductions were observed in acyanotic (36/305 [11.8%] placebo versus 15/300 [5.0%] Synagis) and cyanotic children (27/343 [7.9%] placebo versus 19/339 [5.6%] Synagis).

The clinical studies do not suggest that RSV infection was less severe among children hospitalized with RSV infection who received Synagis for RSV prophylaxis compared to those who received placebo.

## 16 HOW SUPPLIED/STORAGE AND HANDLING

Synagis is supplied in single-dose vials as a preservative-free, sterile liquid solution at 100 mg per mL for intramuscular injection.

50 mg vial NDC 60574-4114-1

The 50 mg vial contains 50 mg Synagis in 0.5 mL.

100 mg vial NDC 60574-4113-1

The 100 mg vial contains 100 mg Synagis in 1 mL.

The rubber stopper used for sealing vials of Synagis is not made with natural rubber latex.

### Storage

Upon receipt and until use, Synagis should be stored between 2°C and 8°C (36°F and 46°F) in its original container. DO NOT freeze. DO NOT use beyond the expiration date.

## 17 PATIENT COUNSELING INFORMATION

### o Advise the patient to read the FDA-approved patient labeling (Patient Information)

The healthcare provider should discuss the potential benefits and risks of Synagis with the parents or guardians of Synagis recipients. Parents or guardians should be informed of the possible side effects of Synagis and of the signs and symptoms of potential allergic reactions and should be advised of the appropriate actions. Parents or guardians should understand the dosing schedule and the importance of compliance with the full course of therapy.

Synagis® is a registered trademark of MedImmune, LLC.



### Manufactured by:

MedImmune, LLC  
Gaithersburg, MD 20878  
U.S. License No. 1799  
1-877-633-4411

RAL-SYNNV17  
Component No.: 26920A

**PATIENT INFORMATION**  
**SYNAGIS®** (Sĭ-nă-jĭs)  
(palivizumab)  
Injection

Read this Patient Information before your child starts receiving SYNAGIS and before each injection. The information may have changed. This leaflet does not take the place of talking with your child's healthcare provider about your child's condition or treatment.

### What is SYNAGIS?

SYNAGIS is a prescription medication that is used to help prevent a serious lung disease caused by Respiratory Syncytial Virus (RSV). Your child is prescribed SYNAGIS because he or she is at high risk for severe lung disease from RSV.

SYNAGIS contains man-made, disease-fighting proteins called antibodies. These antibodies help prevent RSV disease. Children at high risk for severe RSV disease often do not have enough of their own antibodies. SYNAGIS is used in certain groups of children to help prevent severe RSV disease by increasing protective RSV antibodies.

SYNAGIS is not used to treat the symptoms of RSV disease once a child already has it. It is only used to prevent RSV disease.

SYNAGIS is not for adults or for children older than 24 months of age at the start of dosing.

### Who should not receive SYNAGIS?

Your child should not receive SYNAGIS if they have ever had a severe allergic reaction to it. Signs and symptoms of a severe allergic reaction could include:

- severe rash, hives, or itching skin
- swelling of the lips, tongue, or face
- closing of the throat, difficulty swallowing
- difficult, rapid, or irregular breathing
- bluish color of skin, lips, or under fingernails
- muscle weakness or floppiness
- a drop in blood pressure
- unresponsiveness

### What should I tell my child's healthcare provider before my child receives SYNAGIS?

#### Tell your child's healthcare provider about:

- **any reactions** you believe your child has ever had to SYNAGIS.
- **any bleeding or bruising problems.** SYNAGIS is given by injection. If your child has a problem with bleeding or bruises easily, an injection could cause a problem.
- **any other medical problems.**

**Tell your child's healthcare provider about all the medicines your child takes, including prescription and non-prescription medicines, vitamins, and herbal supplements.** Especially tell your child's healthcare provider if your child takes a blood thinner medicine.

### How is SYNAGIS given?

- SYNAGIS is given as a monthly injection, usually in the thigh (leg) muscle, by your child's healthcare provider. Your child's healthcare provider will prescribe the amount of SYNAGIS that is right for your child (based on their weight).
- Your child's healthcare provider will give you detailed instructions on when SYNAGIS will be given.
  - "RSV season" is a term used to describe the time of year when RSV infections most commonly occur (usually fall through spring in most parts of the country). During this time, when RSV is most active, your child will need to receive SYNAGIS shots. Your child's healthcare provider can tell you when the RSV season starts in your area.
  - Your child should receive their **first SYNAGIS shot before the RSV season starts** to help protect them before RSV becomes active. If the season has already started, your child should receive their first SYNAGIS shot as soon as possible to help protect them when exposure to the virus is more likely.

- **SYNAGIS is needed every 28-30 days during the RSV season.** Each dose of SYNAGIS helps protect your child from severe RSV disease for about a month. **Keep all appointments with your child's healthcare provider.**

- **If your child misses an injection, talk to your healthcare provider and schedule another injection as soon as possible.**
- Your child may still get severe RSV disease after receiving SYNAGIS; talk to your child's healthcare provider about what symptoms to look for. If your child has an RSV infection, they should continue to get their scheduled SYNAGIS injections to help prevent severe disease from new RSV infections.
- If your child has certain types of heart disease and has corrective surgery, your healthcare provider may need to give your child an additional SYNAGIS injection soon after surgery.

### What are the possible side effects of SYNAGIS?

#### **Synagis may cause serious side effects including:**

- Severe allergic reactions (may occur after any dose of SYNAGIS). Such reactions may be life-threatening or cause death.
  - See "Who should not take SYNAGIS?" for a list of signs and symptoms.
- Unusual bruising or groups of tiny red spots on the skin.

**Call your child's healthcare provider or get medical help right away if your child has any of the serious side effects listed above after any dose of SYNAGIS.**

#### **Common side effects of SYNAGIS include:**

- fever
- rash

Other possible side effects include skin reactions around the area where the shot was given (like redness, swelling, warmth, or discomfort).

These are not all the possible side effects of SYNAGIS. Tell your child's healthcare provider about any side effect that bothers your child or that does not go away.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

You may also report side effects to MedImmune at 1-877-633-4411.

### General Information about SYNAGIS

Medicines are sometimes prescribed for purposes other than those listed in Patient Information leaflets.

This leaflet summarizes important information about SYNAGIS. If you would like more information, talk with your healthcare provider. You can ask your pharmacist or healthcare provider for information about SYNAGIS that is written for health professionals.

For more information, go to [www.synagis.com](http://www.synagis.com) or call 1-877-633-4411.

### What are the ingredients in SYNAGIS?

Active Ingredient: palivizumab

Inactive Ingredients: chloride, glycine, and histidine

### What is RSV?

Respiratory Syncytial Virus (RSV) is a common virus that is easily spread from person to person. RSV infects nearly all children by their second birthday. In most children, RSV infection is usually no worse than a bad cold. For some children, RSV infection can cause serious lung disease (like pneumonia and bronchiolitis) or breathing problems, and affected children may need to be admitted to the hospital or need emergency care.

Children who are more likely to get severe RSV disease (high-risk children) include babies born prematurely (35 weeks or less) or babies born with certain heart or lung problems.

This Patient Information has been approved by the U.S. Food and Drug Administration.

Synagis® is a registered trademark of MedImmune, LLC.

 **MedImmune**

Manufactured by: MedImmune, LLC

Gaithersburg, MD 20878

Issued March 2014